

The opioid epidemic impacts your patients.

Rates of hospitalizations due to opioid use disorder (OUD) nearly doubled from 2000-2012

Methadone and buprenorphine prevent painful withdrawal and overwhelming cravings in the hospital, which allows the treatment of other medical issues

Starting methadone or buprenorphine can decrease opioid use, HIV, infections, and criminal behavior

Patients with OUD who are not on methadone and buprenorphine have 3x the mortality rate of patients on these medications

We have treatments that can help!

SHOUT

SUPPORT FOR HOSPITAL
OPIOID USE TREATMENT

SHOUT is a coalition of hospitalists and pregnancy care providers working to improve care for patients with opioid use disorder.

The project offers
free resources and support
to launch buprenorphine and methadone at your hospital.

- TOOLKIT FOR IMPLEMENTATION
- SKILL-BUILDING WEBINAR SERIES
- EVIDENCE-BASED GUIDELINES FOR INDUCTION, MAINTENANCE, AND PAIN MANAGEMENT
 - GRAND ROUNDS LED BY AN EXPERT TO BUILD KNOWLEDGE AND MOTIVATION
 - COACHING SESSIONS TO GUIDE YOUR TEAM TOWARDS SUCCESS

FOR MORE INFORMATION OR TO JOIN THE COALITION

Visit the website: <http://tinyurl.com/SHOUTsignup> OR Email: project.SHOUT.coalition@gmail.com



California
Health Care
Foundation



University of California
San Francisco